THE

BOTANICAL EXCHANGE CLUB.

LIST OF DESIDERATA

FOR 1876.

LONDON:

RANKEN AND CO., PRINTERS, DRURY HOUSE, ST. MARY-LE-STRAND.

1876.



REGULATIONS OF THE CLUB.

- 1. The object of the Botanical Exchange Club is to facilitate the exchange of dried specimens of British plants, especially of critical species and varieties. Anyone wishing to become a member will be admitted on payment to the Secretary of an annual subscription of Five Shillings, and on sending a parcel of dried plants in accordance with the subsequent rules. He will then be entitled to share in the distribution of specimens made in the early part of the year following that in which his subscription and parcel were sent.
- 2. Specimens sent for distribution must be carefully dried; must not exceed in size half-a-sheet of demy (16 by 10 inches); and must illustrate the species they represent as completely as possible. Plants more than sixteen inehes long should be once or twice folded, if by so doing the roots can be preserved. (In the Cyperacea, Gramina, and smaller Ferns no specimens should be sent without roots, except in the case of very rare species, which might be in danger of extermination.) No plant must be sent that is not included in the list of desiderata for the current year, unless it be additional to those enumerated in the 7th edition of the "London Catalogue of British Plants," or be from an unrecorded station, or be an unrecorded variety. But up to the date at which the list of desiderata for the year is issued, the desiderata list of the preceding year remains in force, so that the plants collected before the appearance of the annual desiderata list will be received by the Club if they have been mentioned in the desiderata list of the previous year.

- 3. Each specimen must have a label, bearing the number and name of the species as given in the 7th edition of the "London Catalogue"; also the locality and county where, and the date when, the specimen was collected, and the collector's name. The label, should be affixed to the specimen by cutting a transverse slit in the base of the label through which the specimen may be pushed. Any facts connected with a species which the sender thinks important and suitable for the "Report" should be communicated on a separate piece of paper, written on one side only. Such piece of paper should be about seven inches wide, and the notes on each species should be dated and signed by the writer. Specimens sent not in accordance with Rules 2 and 3 will be destroyed.
- 4. Each parcel should be accompanied by a list of the plants the member wishes to receive from the Club. This list is to be made by drawing a short line before their names in the 7th edition of the "London Catalogue."* Species especially wished for should have a! before the horizontal line. The name of the member and address to which the return parcel is to be sent, and any special directions as to how the parcel is to be sent, should be written on the outside of the Catalogue. When the same copy of the Catalogue is used a second time, the species which are no longer wanted, but which were marked the first time, should have a perpendicular line drawn through the horizontal line. Manuscript lists of desiderata will not be received. Rubi and Rosæ will be sent out in sets only; so, in these genera, the horizontal line should be placed before the name of the genus in the Catalogue.
- 5. Parcels may be sent (carriage paid) to any of the following gentlemen not later than the 31st of December:—Dr. Boswell, Balmuto, Kirkcaldy, Fife; Mr. T. R. A. Briggs, 4, Portland Villas, Plymouth; Mr. J. G. Baker, 25, Sydney Villas, Richmond, London; and Mr. Charles Bailey, care of Ralli Brothers, Peter Street, Manchester. Those that arrive after that date will not be available for the distribution for the current year. Members sending more valuable parcels will have their return parcels selected before those who send inferior ones.
- 6. Members who do not send parcels, but who have been members of the Botanical Exchange Club previous to 1872, and paid their

^{*} Published by Mr. R. Hardwicke, 192, Piccadilly, London, W. (Price 6d., post-free 7d.)

subscription for the current year before December 31st, will share in the annual distribution; but the specimens for such non-contributing members will not be selected until the parcels of all the contributing members have been made up, and they cannot be selected according to lists of desiderata; but *general* instructions, such as the desire to receive plants of particular districts, natural orders, or genera, will be attended to by the distributor. In future no new members will be received by the Botanical Exchange Club, except those who contribute specimens of plants.

LIST OF DESIDERATA FOR 1876.

N.B.—Of any of the species in the following List, specimens varying in number from 10 to 50, according to its degree of rarity, will be acceptable.

```
Thalictrum montanum.
            flexuosum.
           majus, var. a.
            Kochii.
           saxatile.
           sphærocarpum.
           riparium
           Morisonii.
           nigricans.
Anemone apennina.
          ranunculoides.
Adonis autumnalis.
Ranunculus Bachii.
            truncatus.
            floribundus.
            penicillatus.
            elongatus.
            radians.
            Godronii.
            confusus.
            marinus.
            salsuginosus.
            intermedius.
            ophioglossifolius.
            reptans.
            vulgatus.
            tomophyllus.
            parvulus.
            chærophyllus.
Caltha minor.
       radicans.
Eranthis hyemalis.
Delphinium Ajacis.
            Consolida.
Pæonia corallina.
Epimedium alpinum.
Nuphar intermedia.
```

pumila.

```
Papaver somniferum.
Glaucium violaceum.
phœniceum.
Chelidonium laciniatum (sponte).
Fumaria Boræi.
          muralis.
         densiflora.
Crambe maritima.
Raphanus maritimus.
Sinapis incana.
Brassica Napus.
         Rutabaga.
         sylvestris.
         Briggsii.
         Cheiranthus.
Hesperis matronalis.
Matthiola sinuata.
          incana.
Dentaria bulbifera.
Arabis petræa.
       ciliata.
       hispida.
       Turrita.
Barbarea arcuata
         intermedia.
Nasturtium siifolium.
Cochlearia Hortii.
           gemina.
Draba inflata.
      muralis.
      aizoides.
Camelina sativa.
         sylvestris.
Thlaspi sylvestre.
Lepidium sativum (sub-sponte).
Helianthemum Breweri.
                vineale.
                grandiflorum.
```

771.	
Viola permixta.	Medicago apiculata.
sepincola.	minima.
calcarea.	
arenaria.	lappacea. Trifolium suffocatum.
lactea.	Townsendi.
intermedia.	
stagnina.	filiforme (Scotland). Oxytropis campestris.
Curtisii, vars. a, b, et c.	Astragalus alpinus.
amœna.	Ornithonus obviortes
Drosera obovata.	Ornithopus ebracteatus. Vicia gracilis.
Polygala grandiflora.	Orobus (fmit)
austriaca.	Orobus (fruit). Lathyrus tuberosus.
uliginosa.	polyetria (f:4)
Dianthus glaucus.	palustris (fruit). acutifolius.
Caryophyllus.	sphæricus.
Silene Otites.	Orobus niger.
italica.	Aremonia agrimonioides.
Lychnis alpina (England).	Prunus insititia.
Holosteum umbellatum.	domestica.
Cerastium pumilum.	Alchemilla conjuncta (sponte).
holosteoides.	Potentilla procumbens.
pentandrum.	norvegica.
alpestre.	Fragaria elatior (sub-sponte).
pubescens.	Rubus fruticosus (any forms named by
compactum.	competent authority).
nigrescens.	Chamæmorus (Wales).
Andrewsii.	Dryas depressa.
Stellaria Boræana.	Rosa (must be named by competent
neglecta.	authority).
Arenaria norvegica.	rubella.
ciliata.	involuta, vars.
rubella.	hibernica.
hybrida.	cordifolia.
uliginosa. Sagina alpina.	pomifera.
nivalis.	mollissima, vars.
Spergularia salina.	tomentosa, vars.
media.	rubiginosa, vars, hand o
Herniaria glabra (Eastern counties).	micrantha, vars.
Scleranthus biennis.	sepium, vars.
perennis.	canina (Bakerianæ).
Claytonia alsinoides.	stylosa, vars.
Elatine Hydropiper (fruit).	austriaca.
Hypericum angustifolium.	Cotoneaster vulgaris.
linarifolium.	Mespilus germanica.
hircinum (sub-sponte).	Fyrus latifolia.
elatum (sub-sponte).	scandica.
Althæa hirsuta.	semipinnata.
Malva borealis.	fennica.
parviflora, L.	Pyraster.
Tilia grandifolia (sponte).	Achras.
parvifolia.	Briggsii.
Linum perenne.	Lythrum Hyssopifolia.
Geranium prostratum.	Enothera odorata.
lancastriense (sponte).	Isnardia palustris.
modestum.	Circæa intermedia (lutetiana).
purpureum.	Intermedia (alnina)
striatum (sub-snonte)	bi yriophyllum pectinatum
Erodium chærophyllum	Camtriche verna (fruit).
Ulex nanus.	obtusangula.
Lupinus perennis.	truncata.
Ononis repens.	Ribes sylvestre.
reclinata	Smithianum.
	spicatum.

Sedum teretifolium.	Bidens radiata.
micranthum.	hybrida.
dasyphyllum.	Inula salicina.
sexangulare.	Aster salignus.
albescens.	Petasites fragrans.
rupestre.	Hypochæris Balbisii.
Forsterianum.	Picris arvalis.
Sempervivum tectorum (quasi-sponte).	Tragopogon grandiflorus.
Saxifraga stellaris, var. vivipara.	porrifolius (sub-sponte).
Geum, vars. (Irish).	Taraxacum lævigatum.
hirsuta.	Lactuca Scariola.
umbrosa.	Mulgedium alpinum.
cernua.	Sonchus glabra.
cæspitosa.	palustris.
Sternbergii.	Crepis fœtida.
decipiens.	taraxacifolia.
sponhemica.	nicæensis.
Astrantia major.	Hieracium. (Any named by competen
Eryngium campestre (Kent).	authority, except —
Helosciadium repens.	Pilosella.
Moorei	murorum.
Trinia vulgaris.	vulgatum (type).
Bupleurum aristatum (Sussex).	umbellatum.
Enanthe fluviatilis.	boreale.
Siler trilobum.	Phyteuma spicatum.
Meum Athamanticum.	Campanula latifolia purpurea.
Peucedanum palustre.	persicifolia (sub-sponte).
Tordylium maximum.	Vaccinium uliginosum.
Caucalis latifolia.	Arctostaphylos alpina.
Viscum album (male).	Uva-ursi.
Lonicera Caprifolium.	Arbutus Unedo (sponte).
Linnæa borealis.	Loiseleuria procumbens.
Galium erectum.	Menziesia polifolia.
Mollugo, vars. b and c.	cærulea.
nitidulum.	Erica hibernica.
elongatum.	Pyrola rotundifolia.
anglicum.	uniflora.
Asperula arvensis.	Monotropa hirsuta.
Valerianella carinata.	Erythræa latifolia.
mixta.	Cicendia pusilla.
eriocarna.	Gentiana nivalis.
Carduus litigiosus.	germanica.
tuberosus.	Cuscuta Epilinum.
setosus.	Solanum miniatum.
hybridi (any).	Verbascum pulverulentum.
Arctium intermedium.	Blattaria.
nemorosum.	hybrida (any).
Centaurea aspera.	Linaria Pelisseriana.
solstitialis.	repens, var. b.
Matricaria inodora, vars. b and c.	speciosa.
Anthemis anglica.	purpurea.
Diotis maritima (England and Wales).	Veronica triphylla.
Artemisia campestris.	verna.
Filago apiculata.	saxatilis.
Gnaphalium pilulare.	spicata.
luteo album.	hirsuta.
norvegicum.	tenella.
hyperboreum.	Euphrasia officinalis, vars.
Senecio vulgaris, var. hibernica.	Bartsia rotundata.
paludosus.	Rhinanthus major.
palustris.	Melampyrum ericetorum.
maritima.	Orobanche arenaria.
Doronicum plantagineum	onrilos

Orobanche caryophyllacea.	Chenopodium intermedium.
elatior.	botryoides.
Picridis.	glaucum.
amethystea.	Atriplex erecta, vars. a and b.
Mentha alopecuroides (sponte).	prostrata.
sylvestris, vars. b and c.	Smithii.
vulgaris.	triangularis.
pubescens.	
citrata (quasi-sponte).	Rumov conglomoratus faces f
hirsuta, var. b.	Rumex conglomeratus, forms of.
	rupestris.
sativa, var. subglabra, Baker.	viridis, forms of.
gracilis.	Warrenii.
cardiaca (quasi-sponte),	palustris.
pratensis.	pulcher.
gentilis, vars. b and c.	obtusifolius, forms of.
arvensis, vars. b, c, d, e, f.	elongatus.
erecta.	trigranulatus.
Thymus Serpyllum, forms of.	aquaticus.
Chamædrys.	maximus.
Origanum macrostachyum.	alpinus.
Calamintha Briggsii.	hybrida (any).
Salvia clandestina.	Polygonum a constitution
pratensis.	Polygonum agrestinum.
Ballota ruderalis.	microspermum.
	rurivagum.
Stachys germanica.	littorale.
Galeopsis intermedium.	maritimum.
ochroleuca.	elatum.
Lamium intermedium.	maculatum.
decipiens.	viviparum.
Ajuga pyramidalis.	Daphne Mezereum (leaves).
Teucrium Botrys (Sanderstead).	Asarum europæum.
Scordium.	Aristolochia Clematitis.
Pulmonaria angustifolia.	Euphorbia Peplis.
Lithospermum purpureo-cærulcum.	
Myosotis strigulosa.	stricta.
alpestris (England).	hiberna.
umbrosa.	pseudo-Cyparissias.
Mittenii.	Cyparissias.
Anchusa officinalis.	coralloides.
	dulcis.
Asperugo procumbens.	Mercurialis ambigua.
Pinguicula grandiflora.	Parietaria fallax.
alpina.	Urtica pilulifera.
Utricularia neglecta (flowers).	Dodartii.
minor (flowers).	Ulmus glabra.
intermedia (flowers).	montana, vars. b and c.
Primula elatior.	Quercus intermedia.
scotica, var. b.	Betula alba yang a h
Cyclamen hederifolium.	Betula alba, vars. a, b, and c.
Lysimachia thyrsiflora.	Populus alba (female flowers).
punctata.	canescens (female flowers
ciliata.	tremula, vars, a and h
Armeria duriuscula.	Saux decipiens.
Statice pyramidalis.	viridis.
Dodartii.	Russelliana.
	Woolgariana.
caspia.	Lambertiana.
Plantago hirsuta.	Doniana.
arenaria.	Helix.
Amaranthus Blitum.	intricata.
Salicornia procumbens.	
radicans.	stipularis.
Chenopodium cymosum.	pseudo-stipularis.
ficifolium.	rugosa.
urbicum.	acuminata.
ar oroum.	oleifolia.

Sanx sphacerata.	Epipogon apityrium.
laurina.	Corallorhiza innata.
phylicifolia, vars. named.	Liparis Loeselii.
nigricans, vars. named.	Malaxis paludosa.
ambigua and vars.	Cypripedium Calceolus.
repens, vars. named.	Sisyrinchium Bermudiana.
rosmarinifolia.	Trichonema Columnie.
angustifolia.	Gladiolus illyricus.
lanata.	Iris citrina.
Lapponum and vars.	Bastardi.
	tuberosa.
Arbuscula and vars.	Crocus aureus.
Myrsinites, vars.	biflorus.
Grahami.	1
Bakeri.	Narcissus Bromfieldii.
Moorei.	major.
Sadleri.	pœticus.
Pinus sylvestris (sponte).	Tazetta.
pinaster.	Leucojum æstivum.
Juniperus nana.	vernum.
Taxus fastigiata.	Smilacina bifolia.
Typha media.	Polygonatum verticillatum.
Sparganium affine.	officinale.
Arum italicum.	intermedium.
Potamogeton pseudo-fluitans.	Ruscus aculeatus (in fruit).
linearis.	Lloydia serotina.
	Ornithogalum nutans.
plantagineus.	Muscari racemosum.
sparganiifolius.	
Lonchitis.	Comosum.
nitens.	Allium Ampeloprasum, var. a.
acuminatus.	Babingtonii.
decipions (flowers).	sphærocephalum.
longifolius.	vineale, var. a.
zosterifolius.	complanatum.
acutifolius.	carinatum.
tenuissimus.	paradoxum.
trichoides.	Moly.
pectinatus (typc).	Simethis bicolor.
flabellatus (fruit).	Luzula sudetica.
Zannichellia (any).	Juneus castaneus.
Ruppia spiralis.	filiformis (Scotland).
Zostera nana.	macrocephalus.
Naias flexilis.	Kochii.
Zcheuchzeria palustris.	Rhyncospora sordida.
Sagittaria sagittifolia (winter bulbs).	Scirpis uniglumis.
Alisma lanceolatum.	parvulus.
natans.	Holoschænus (root-leaves)
Stratiotes aloides (male flowers and	pungens.
fruit).	Eriophorum alpinum.
Orchis hircina.	gracile.
purpurea.	Kobresia caricina (England).
militaris.	Carex incurva.
Simia.	divisa.
laxiflora.	pseudo-divulsa.
incarnata.	Boenninghauseniana.
Neotinea intacta.	elongata.
Ophrys aranifera.	lagopina.
fucifera.	alpicola.
arachnites.	bracteata.
Spiranthes æstivalis.	alpina.
gemmipara.	Buxbaumii.
Epipactis ovalis.	stricta.
violacea.	aquatilis.
Cephalanthera rubra.	
oopimanuncia rupra.	Watsoni.

Carex uliginosa.	Triticum biflorum.
Gibșonii.	pungens.
Micheliana.	acutum.
strictocarpa.	Lolium arvense.
irrigua.	linicola.
ornithopoda.	Elymus arenarius.
ericetorum.	Hordeum sylvaticum.
tomentosa.	Hymonophyllum tunbaidana
ustulata.	Hymenophyllum tunbridgense.
	Trichomanes radicans.
frigida.	Andrewsii.
punctata.	Adiantum Capillus-Vencris.
speirostachya.	Asplenium germanicum.
ebracteata.	anceps.
involuta.	lanceolatum.
pulla.	microdon.
Grahami.	acutum.
Chamagrostis minima.	serpentini.
Digitaria humifusa.	fontanum.
Setaria verticillata.	Athyrium molle.
Hierochloa borealis (with root-leaves).	incisum.
Anthoxanthum Puclii.	Woodsie hamenhaue
Alopecurus pronus.	Woodsia hyperborea.
alpinus.	ilvensis.
Watsoni.	Cystopteris fragilis.
	dentata.
Phleum Boehmeri.	Dickieana.
Sesleria flavescens.	alpina.
Lagurus ovatus.	montana.
Polypogon littoralis.	Aspidium Lonchitis (England or Ire-
Agrostis interrupta (England).	land).
Calamagrostis Hookeri.	Nephrodium abbreviatum.
tenella.	pumilum.
Corynephorus canescens.	cristatum.
Aira montana.	remotum.
Avena fatua, vars. a, b, and c.	
Koeleria cristata, vars. b and c.	uliginosum.
Poa stricta.	spinulosum.
alaxa.	collinum.
glauca.	nanum.
Balfourii.	glandulosum.
Parnellii.	Polypodium alpestre.
Cynosurus echinatus.	flexile.
Festuca ambigua.	Gymnogramme leptophylla.
	Ophioglossum ambiguum.
glauca.	lusitanicum.
major.	Botrychium rutaceum.
arenaria.	Lycopodium annotinum.
arundinacea.	Isoetes echinospora.
Bromus Benekenii.	Hystrix.
midritensis.	Equisetum pratense (fruct.).
rigidus.	alnostro
maximus.	alpestre.
pubescens.	subnudum.
Lloydianus.	paleaceum.
patulus.	Wilsoni.
Brachypodium glabrescens.	trachyodon.
pubosses s	Chara (any named by competent
pubescens.	authority).
	J /

